

Control of Crabgrass in Home Lawns

Purdue University Turf Science

Department of
Agronomy
www.agry.purdue.edu/turf

University of Illinois Turfgrass Program

Department of
Natural
Resources and
Environmental
Sciences
www.turf.uiuc.edu

Crabgrass is a common weed that infests home lawns in the Midwest (Figure 1). Crabgrass is an summer annual weed that germinates when soil temperatures are approximately 60° F for 3-5 days at the 1/4" level. It begins flowering and setting seed in July and dyes with the first frost of fall. Crabgrass has tremendous survival reproductive capabilities. Because of this, it is unrealistic to expect a crabgrass free lawn. You cannot eradicate crabgrass (or any other pest for that matter); a few crabgrass plants in your lawn are acceptable.

Cultural Crabgrass Control

The most effective way to control crabgrass is to create a dense, healthy turf. A healthy turf will compete well with crabgrass and prevent it from establishing.

Mowing

- Mow at 2.5 to 3.0 inches depending on the turf species. Mowing below this range will increase crabgrass populations (Figure 2).
- Mow frequently so as not to remove more than 1/3 of the leaf blade at one time. This may mean mowing twice weekly in spring and every other week in summer.

Irrigation

Irrigate deeply and infrequently. Daily, light irrigations promote shallow rooting, non-drought hardy turf, and encourage crabgrass. Water to wet the soil to the depth of rooting, and then do not water again until you see the first sign of drought stress (When drought stressed, turf will become bluish gray and footprints will remain in the turf after it is walked on).



Figure 1. Mature crabgrass plant (Zac Reicher).




Figure 2. High crabgrass population in a Kentucky bluegrass lawn mown at 1.0 inch (Aaron Patton).

Fertilization

Apply 2 to 4 pounds nitrogen per 1000 ft² each year to create a dense lawn and reduce crabgrass populations. Apply 60-100% of the nitrogen in two applications in Fall: one in September and one in November after the final mowing. Avoid applications of nitrogen in summer that increase crabgrass vigor.

Chemical Control

Often, cultural control alone will not control crabgrass satisfactorily, and herbicides may be needed. This is especially true in new lawns or lawns that are thin from damage or improper maintenance. When using herbicides and all pesticides, be sure to read, understand, and follow all label recommendations.

Preemergence Herbicides

Preemergence herbicides prevent emergence of crabgrass plants. These products must be applied prior to crabgrass emergence which could occur as early as April 1 in southern Indiana and Illinois and three or more weeks later in northern Indiana and Illinois (Figure 3). Purdue research has shown that these herbicides can be applied as early as March 1 and still be effective all season. It is essential to apply these products early in spring prior to crabgrass germination.

Often, preemergence herbicides are combined with fertilizers as weed and feed products. Since fertilization should be minimized in the spring, purchase products with most of the nitrogen in slow release forms such as methylene ureas or sulfur or polymer-coated ureas. Avoid products with mostly quick release nitrogen such as urea or ammonical nitrogen.

Do not use preemergence herbicides on new seedlings or before seeding an area. To be most effective, these products need to be watered-in after application. Refer to the label for specific instructions of each product.

Common Names of Preemergence Herbicides

Benefin
Oxadiazon
Benefin/Trifluralin
Pendimethalin
Dithiopyr
Prodiamine
Corn Gluten

Postemergence Herbicides

Postemergence herbicides control crabgrass after it has emerged and are most effective on small crabgrass plants. These products are more difficult to use than preemergence herbicides and it is extremely important to follow label instructions. Of the products listed below, quinclorac is safest for turfgrass seedlings. Keep in mind the following when using these products:

- Be sure to read, understand and follow all herbicide label directions for the safest, most effective weed control.
- The area must be well-watered prior to application and not under drought stress.
- Do not mow or water for 24 hours following application.
- Apply at temperatures below 85° F. These products are most effective on clear days with low humidity.
- A second application may be needed within seven days for most effective control.
- Refer to the label for use before and after seeding.

Common Names of Postemergence Herbicides

MSMA (Monosodium methyl arsonate)
DSMA (Disodium methyl arsonate)
Dithiopyr (only effective on crabgrass seedlings)
Fenoxaprop
Quinclorac

Do not attempt to control crabgrass with herbicides after mid- July because crabgrass plants are usually too large to control effectively. It is better to simply tolerate the crabgrass until it dies with the first frost.

By maintaining a dense lawn, you can limit the amount of crabgrass. Proper fertility, mowing, and irrigation is essential for crabgrass control; consider herbicidal control only if necessary.



Figure 3. Predicted crabgrass germination date in Illinois and Indiana based on normal weather data. Remember that preemergence herbicides must be applied at least two weeks prior to these dates to control crabgrass

Authors:

Zac Reicher, Professor and Turfgrass Extension Specialist, Purdue University Department of Agronomy
Cale Bigelow, Assistant Professor and Turfgrass Extension Specialist, Purdue University Department of Agronomy
Aaron Patton, Graduate Research Assistant, Purdue University Department of Agronomy
Tom Voigt, Associate Professor and Turfgrass Extension Specialist, University of Illinois Department of Natural Resources and Environmental Sciences



Rev. 5/2006

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Order or download materials on this and other topics from:

Purdue Extension Education Store: www.ces.purdue.edu/new
University of Illinois Extension Publications Plus: www.pubsplus.uiuc.edu